

HARDER-NARASIMHAN FILTRATIONS AND K-GROUPS OF AN ELLIPTIC CURVE

GUODONG ZHOU

ABSTRACT. Let X be an elliptic curve over an algebraically closed field. We prove that some exact sub-categories of the category of all vector bundles over X , defined using Harder-Narasimhan filtrations, have the same K-groups as the whole category.

1. INTRODUCTION

Throughout this paper, k denotes an algebraically closed field. Let X be a smooth projective curve over k and let E be a vector bundle over X . We define the slope of E as the quotient of its degree by its rank, i.e. $\mu(E) = \deg(E)/\text{rank}(E)$. A vector bundle E is called semi-stable (resp. stable) if for any non-zero proper sub-bundle E' , we have $\mu(E') \leq \mu(E)$ (resp. $\mu(E') < \mu(E)$). The importance of the notion of semi-stability consists in the constructions of moduli spaces of vector bundles, see for example [8][12][13][6][4]. For each vector bundle E , there exists a unique filtration, say *Harder-Narasimhan filtration* ([5, Proposition 1.3.9]),

$$0 = E_0 \subsetneq E_1 \subsetneq \cdots \subsetneq E_{s-1} \subsetneq E_s = E$$

such that the quotients $F_i = E_i/E_{i-1}$ are semi-stable for all $1 \leq i \leq s$ and

$$\mu(F_1) > \mu(F_2) > \cdots > \mu(F_s).$$

We note $\mu_{\max}(E) = \mu(F_1)$ and $\mu_{\min}(E) = \mu(F_s)$.

Let $\mathcal{P}(X)$ be the exact category of all vector bundles over X . Let $I \subset \mathbb{R}$ be a connected interval (possibly of length zero). Following T. Bridgeland ([2, Section 3]), denote by $\mathcal{P}(I)$ the full sub-category of $\mathcal{P}(X)$ consisting of all vector bundles E such that $\mu_{\max}(E), \mu_{\min}(E) \in I$. It is an interesting fact that the category $\mathcal{P}(I)$ is also exact with the exact category structure induced from that of $\mathcal{P}(X)$ (see Lemma 2.1 below). We can therefore consider K-groups of $\mathcal{P}(I)$, as defined by D.Quillen for an exact category using his famous Q -construction ([11]). In this paper, we are interested in the relations between K-groups of $\mathcal{P}(I)$ and K-groups of $\mathcal{P}(X)$, i.e. those of X in case that X is an elliptic curve. More precisely, we prove the following theorem.

Theorem 1.1. *Let X be an elliptic curve over k an algebraically closed field and let I be a connected interval of strictly positive length. Then the inclusion functor $\mathcal{P}(I) \hookrightarrow \mathcal{P}(X)$ induces isomorphisms of K-groups $K_i(\mathcal{P}(I)) \xrightarrow{\cong} K_i(X)$ for all $i \geq 0$.*

Vector bundles over an elliptic curve were classified by M.Atiyah in [1]. His classification is essential to the proof of the preceding theorem. Roughly speaking, the idea is to construct, for an enough general vector bundle, a resolution of length one in $\mathcal{P}(I)$ and then the resolution theorem ([11, Theorem 3.3]) applies.

The following question is natural.

2000 Mathematics Subject Classification: 19E08, 14H52, 14H60
Key words: elliptic curve, Harder-Narasimhan filtration, K-groups, the resolution theorem, semi-stable vector bundles.

Question 1.2. *Does the statement in the preceding theorem hold if we replace X by any smooth projective curve of genus ≥ 2 ?*

Acknowledgement: I want to express my gratitude to Mr. Bruno Kahn (Directeur de Recherche du CNRS) for some useful conversations.

2. PROOF OF THE MAIN THEOREM

Firstly we prove the following fact mentioned in the introduction.

Lemma 2.1. *The category $\mathcal{P}(I)$ is an exact category whose exact sequences are given by short exact sequences in $\mathcal{P}(X)$ with their terms in $\mathcal{P}(I)$.*

Proof. One needs to show that $\mathcal{P}(I)$ is closed under extensions. Take a short exact sequence

$$0 \rightarrow E' \rightarrow E \rightarrow E'' \rightarrow 0$$

with $E', E'' \in \mathcal{P}(I)$. Let

$$0 = E_0 \subsetneq E_1 \subsetneq \cdots \subsetneq E_{s-1} \subsetneq E_s = E$$

be the Harder-Narasimhan filtration of E . We then have the exact sequence

$$0 \rightarrow E' \cap E_{s-1} \rightarrow E' \rightarrow F \rightarrow 0$$

with F a sub-bundle of $F_{s-1} = E/E_{s-1}$. We obtain that

$$\mu_{\min}(E) = \mu(F_{s-1}) \geq \mu(F) \geq \mu_{\min}(E').$$

We also have the exact sequence

$$0 \rightarrow E' \cap E_1 \rightarrow E_1 \rightarrow G \rightarrow 0$$

with G a sub-bundle of E'' . We get that $\mu(E' \cap E_1) \leq \mu_{\max}(E')$ and $\mu(G) \leq \mu(E'') \leq \mu_{\max}(E'')$ and as $\mu_{\max}(E) = \mu(E_1)$ is the barycenter of $\mu(E' \cap E_1)$ and $\mu(G)$ with positive coefficients, $\mu_{\max}(E) \leq \mu_{\max}(E'), \mu_{\max}(E'')$. This prove that $E \in \mathcal{P}(I)$. □

Next we recall some known facts about vector bundles over an elliptic curve X .

Lemma 2.2. [6, Chapter 8, Section 8.7, Exercise 2.2] *Each vector bundle over X is a direct sum of indecomposable bundles. In particular, every indecomposable vector bundle is semi-stable.*

Theorem 2.3. *Let E and F be two semi-stable vector bundles over an elliptic curve. Then $E \otimes F$ is still semi-stable.*

In fact, in case of characteristic zero, the tensor product of two semi-stable vector bundles is semi-stable over a smooth projective curve of arbitrary genus. This was first proved by M.S.Narasimhan and C.S.Seshadri using analytic method ([9]) and then by Y. Miyaoka using algebraic method ([7, Corollary 3.7]). The case of positive characteristic uses the notion of strong semi-stability. A vector bundle is called strongly semi-stable if all its Frobenius pullbacks are semi-stable. T. Oda proved in [10, Theorem 2.16] (see also [14, Corollary 3^p]) that a semi-stable vector bundle over an elliptic curve is strongly semi-stable. Then the preceding theorem follows from the facts ([7, Section 5]) that the tensor product of two strongly semi-stable vector bundles is still strongly semi-stable and that strong semi-stability implies semi-stability.

Let $\mathcal{E}(r, d)$ with $r \geq 1$ and $d \in \mathbb{Z}$ be the set of isomorphism classes of indecomposable vector bundles of rank r and of degree d . When r and d are coprime, M. Atiyah introduced a distinguished vector bundle $E_{r,d} \in \mathcal{E}(r, d)$ (Atiyah noted it by $E_A(r, d)$) with the property $E_{r,d}^* \cong E_{r,-d}$ ([1, Corollary of Theorem 7]).

Let us construct the resolutions of length one for an enough general vector bundle. The starting point is the following lemma.

Lemma 2.4. *Let $E \in \mathcal{E}(r, d)$ with $r \geq 1$ and $d > 0$. Then there exists a vector bundle $E' \in \mathcal{E}(r + d, d)$, unique up to isomorphisms, given by the extension*

$$0 \rightarrow H^0(E) \otimes \mathcal{O}_X \rightarrow E' \rightarrow E \rightarrow 0$$

Moreover, $H^0(E) \cong H^0(E')$ and the map $H^0(E') \otimes \mathcal{O}_X \cong H^0(E) \otimes \mathcal{O}_X \rightarrow E'$ is the evaluation map.

Proof. The existence of E' follows from [1, Lemma 16] and other statements are easy consequences of [1, Lemma 15]. \square

Proposition 2.5. *Let $E \in \mathcal{E}(r, d)$ with $r \geq 1$ and $d > 0$ and let $\epsilon > 0$. There exists a short exact sequence*

$$0 \rightarrow E_1 \rightarrow E_0 \rightarrow E \rightarrow 0$$

where E_1 is semi-stable of zero slope and where E_0 is semi-stable of slope $\mu(E_0) \in (0, \epsilon)$.

Proof. The preceding lemma gives an exact sequence

$$0 \rightarrow H^0(F_1) \otimes \mathcal{O}_X \xrightarrow{ev} F_1 \xrightarrow{f_1} E \rightarrow 0$$

with $F_1 \in \mathcal{E}(r + d, r)$ and where ev is the evaluation map. We again apply Lemma 2.4 to F_1 and we obtain

$$0 \rightarrow H^0(F_2) \otimes \mathcal{O}_X \xrightarrow{ev} F_2 \xrightarrow{f_2} F_1 \rightarrow 0$$

with $F_2 \in \mathcal{E}(r + 2d, d)$. These two exact sequences yield

$$0 \rightarrow \text{Ker}(f_1 \circ f_2) \rightarrow F_2 \xrightarrow{f_1 \circ f_2} E \rightarrow 0$$

and

$$0 \rightarrow H^0(F_2) \otimes \mathcal{O}_X \rightarrow \text{Ker}(f_1 \circ f_2) \rightarrow H^0(F_1) \otimes \mathcal{O}_X \rightarrow 0 \quad (*)$$

Lemma 2.1 implies that $\text{Ker}(f_1 \circ f_2)$ is semi-stable of zero slope.

If we iterate this process for n times with n enough great such that $d/(r + nd) < \epsilon$, we get

$$0 \rightarrow \text{Ker}(f_1 \circ \cdots \circ f_n) \rightarrow F_n \xrightarrow{f_1 \circ \cdots \circ f_n} E \rightarrow 0 \quad (**).$$

As above, it is easy to show that $\text{Ker}(f_n \circ \cdots \circ f_0)$ is semi-stable of zero slope and that $(**)$ is the desired resolution. \square

Now we give the proof of the main theorem.

Proof. (of Theorem 1.1)

We can suppose that $I = (a, b)$ with $-\infty < a < b < +\infty$. For any real number λ , we note $I + \lambda = (a + \lambda, b + \lambda)$. Set $J = (a, +\infty)$.

Step I: We show that the inclusion functor $\mathcal{P}(I) \hookrightarrow \mathcal{P}(J)$ induces isomorphisms of K -groups. Take two integers $r \geq 1$ and d such that $-\frac{d}{r} = -\mu \in I$, $(r, d) = 1$ and $(r, p) = 1$ if $\text{char } k = p > 0$. By Theorem 2.3, the tensor product by $E_{r,d}$ is an exact functor from $\mathcal{P}(I)$ to $\mathcal{P}(I + \mu)$. Note that $0 \in I + \mu$. Let $E \in \mathcal{P}(J)$. Then $E \otimes E_{r,d} \in \mathcal{P}((a + \mu, +\infty))$. Suppose that $E \otimes E_{r,d} = \oplus F_i$ with all F_i indecomposable. If $F_i \in \mathcal{P}(I + \mu)$, then we take the resolution

$$0 \rightarrow 0 \rightarrow F_i \xrightarrow{Id} F_i \rightarrow 0$$

and if $F_i \notin \mathcal{P}(I + \mu)$, we take the resolution given by Proposition 2.5 with $\epsilon = b + \mu$. The sum of these resolutions of all F_i is a resolution of E of the form

$$0 \rightarrow E_1 \rightarrow E_0 \rightarrow E \otimes E_{r,d} \rightarrow 0$$

where E_1 is semi-stable of zero slope and where E_0 is in $\mathcal{P}(I + \mu)$. Now the tensor product of the resolution above by $E_{r,-d}$ gives

$$0 \rightarrow E_1 \otimes E_{r,-d} \rightarrow E_0 \otimes E_{r,-d} \xrightarrow{f} E \otimes E_{r,d} \otimes E_{r,-d} \rightarrow 0$$

By [10, Corollary 2.7], $E_{r,d} \otimes E_{r,-d} \cong \mathcal{E}nd(E_{r,d}) = \mathcal{O}_X \oplus G$. We write g the projection from $E \otimes E_{r,d} \otimes E_{r,-d}$ to $E \otimes G$. We have an exact sequence

$$0 \rightarrow E_1 \otimes E_{r,-d} \rightarrow \text{Ker}(g \circ f) \rightarrow E \rightarrow 0$$

Obviously $E_1 \otimes E_{r,-d}$ is semi-stable of slope $-\mu$. The inequality $\mu_{\max}(\text{Ker}(g \circ f)) \leq \mu_{\max}(E_0 \otimes E_{r,-d})$ together with Lemma 2.1 implies that $\text{Ker}(g \circ f) \in \mathcal{P}(I)$. The resolution theorem applies and we obtain that the inclusion functor $\mathcal{P}(I) \hookrightarrow \mathcal{P}(J)$ induces isomorphisms of K-groups.

Step II: We show that the inclusion functor $\mathcal{P}(J) \hookrightarrow \mathcal{P}(X)$ induces isomorphisms of K-groups. By a theorem of Serre ([3, Chapter 2, Theorem 5.17]), for each $E \in \mathcal{P}(X)$, we have an exact sequence

$$0 \rightarrow E \rightarrow \mathcal{O}_X(n)^m \rightarrow F \rightarrow 0$$

with $n, m \gg 0$ and then $\mathcal{O}_X(n)^m, F \in \mathcal{P}(J)$. Let us consider the functor $\mathcal{P}(J)^{op} \hookrightarrow \mathcal{P}(X)^{op}$ where op means the opposite category. Notice that $QC^{op} \cong QC$ ([11, Page 94]) where Q is the Q -construction and then $K_i(\mathcal{C}^{op}) \cong K_i(\mathcal{C})$ for all $i \geq 0$, we can deduce from the resolution theorem that the inclusion functor $\mathcal{P}(J) \hookrightarrow \mathcal{P}(X)$ induces isomorphisms of K-groups.

This finishes the proof. □

REFERENCES

- [1] M.F.Atiyah, *Vector bundles over an elliptic curve*, Proc. London Math. Soc. (3) **7** (1957), 414-452
- [2] T.Bridgeland, *Stability conditions on triangulated categories*, Ann. Math. **166** (2007), 317-346
- [3] R.Hartshorne, *Algebraic Geometry*, Graduate Texts in Mathematics **52**, Springer-Verlag 1977
- [4] D.Huybrechts and M.Lehn, *The Geometry of Moduli Spaces of Sheaves on Curves*, Aspect of Mathematics E31, Friedr. Vieweg & Sohn, Braunschweig 1997
- [5] G.Harder and M.S.Narasimhan, *On the cohomology groups of moduli spaces of vector bundles on curves*, Math. Ann. **212** (1974/75), 215-248
- [6] J.Le Potier, *Lecteur on Vector Bundles*, Cambridge Studies in Advanced Mathematics **54**, Cambridge University Press, 1997
- [7] Y.Miyaoka, *The chern class and Kodaria dimension of a minimal variety*, (Algebraic geometry, Sendai, 1995), Advanced Studies in Pure Mathematics **10** (1987), 449-476
- [8] D.Mumford, *Projective invariants of projective structures and applications* Proc. ICM Stockholm (1962) 526-530
- [9] M.S.Narasimhan and C.S.Seshadri, *Stable bundles and unitary bundles on a compact Riemann surface*, Ann. Math. (2) **82** (1965), 540-567
- [10] T.Oda, *Vector bundles on an elliptic curve*, Nagoya Math. J. **43** (1971), 41-72
- [11] D.Quillen, *Higher algebraic K-theory I*, Lect. Notes in Math. 341, 77-139, Springer, Berlin 1973
- [12] C.S.Seshadri, *Fibrés vectoriels sur les courbes algébriques*, Astérisque **96** (1992)
- [13] C.T.Simpson *Moduli of representations of the fundamental Group of a smooth projective variety I*, Publ. Math. I.H.E.S. **79** (1994), 47-129
- [14] N.I.Sheferd-Barron, *Semi-stability and reduction mod p*, Topology **37** (1998), 659-664

GUODONG ZHOU
LAMFA ET CNRS UMR 6140
UNIVERSITÉ DE PICARDIE JULES VERNE
33, RUE ST LEU
80039 AMIENS
FRANCE
E-mail address: `guodong.zhou@u-picardie.fr`